# ABclonal www.abclonal.com

## **SLC25A23 Rabbit pAb**

Catalog No.: A14352

## **Basic Information**

#### **Observed MW**

52kDa

## **Calculated MW**

52kDa

#### Category

Primary antibody

## **Applications**

ELISA, WB, IF/ICC

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

Predicted to enable ATP transmembrane transporter activity. Involved in calcium import into the mitochondrion; positive regulation of mitochondrial calcium ion concentration; and regulation of cellular hyperosmotic salinity response. Located in mitochondrion.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:200

## **Immunogen Information**

**Gene ID**79085
Swiss Prot
Q9BV35

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-160 of human SLC25A23 (NP\_077008.2).

#### **Synonyms**

APC2; MCSC2; SCAMC3; SCaMC-3; SLC25A23

## **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using SLC25A23 Rabbit pAb (A14352) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.



Immunofluorescence analysis of NIH/3T3 cells using SLC25A23 Rabbit pAb (A14352) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.