Leader in Biomolecular Solutions for Life Science

ACSM2A Rabbit pAb

Catalog No.: A15563 1 Publications



Basic Information

Observed MW

64kDa

Calculated MW

64kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Mouse,Rat

Background

This gene encodes a mitochondrial acyl-coenzyme A synthetase that is specific for medium chain fatty acids. These enzymes catalyze fatty acid activation, the first step of fatty acid metabolism, through the transfer of acyl-CoA. These enzymes also participate in the glycine conjugation pathway in the detoxification of xenobiotics such as benzoate and ibuprofen. Expression levels of this gene in the kidney may be correlated with kidney function. This gene and its paralog ACSM2B (Gene ID: 348158), both present on chromosome 16, likely arose from a chromosomal duplication event.

Recommended Dilutions

WB

1:200 - 1:2000

Immunogen Information

Gene ID

Swiss Prot

123876 Q08AH3

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 418-577 of human ACSM2A (NP_001295101.1).

Synonyms

ACSM2; A-923A4.1; ACSM2A

Contact

•

www.abclonal.com

Product Information


Source Rabbit **Isotype** IgG **Purification**Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using ACSM2A Rabbit pAb (A15563) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.