

A15921

Leader in Biomolecular Solutions for Life Science



TUBGCP6 Rabbit pAb

Catalog No.: A15921

Basic Information

Observed MW

250kDa

Calculated MW

200kDa

Category

Mouse Monoclonal Antibody

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

The protein encoded by this gene is part of a large multisubunit complex required for microtubule nucleation at the centrosome.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

Immunogen Information

Gene ID

85378

Swiss Prot

Q96RT7

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 600-900 of human TUBGCP6 (NP_065194.2).

Synonyms

GCP6; GCP-6; MCCRP; MCCRP1; MCPHCR; TUBGCP6

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of various lysates using TUBGCP6 Rabbit pAb (A15921) at 1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.



Immunohistochemistry analysis of TUBGCP6 in paraffin-embedded Human colon carcinoma using TUBGCP6 Rabbit pAb (A15921) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of TUBGCP6 in paraffin-embedded Rat testis using TUBGCP6 Rabbit pAb (A15921) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of TUBGCP6 in paraffin-embedded Rat testis using TUBGCP6 Rabbit pAb (A15921) at dilution of 1:100 (20x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.