Leader in Biomolecular Solutions for Life Science

# **NOSTRIN Rabbit pAb**

Catalog No.: A15938



## **Basic Information**

## **Observed MW**

55kDa

## **Calculated MW**

58kDa

#### Category

Polyclonal Antibody

## **Applications**

WB,IF/ICC,ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

Nitric oxide (NO) is a potent mediator in biologic processes such as neurotransmission, inflammatory response, and vascular homeostasis. NOSTRIN binds the enzyme responsible for NO production, endothelial NO synthase (ENOS; MIM 163729), and triggers the translocation of ENOS from the plasma membrane to vesicle-like subcellular structures, thereby attenuating ENOS-dependent NO production.

## **Recommended Dilutions**

**WB** 1:500 - 1:1000

**IF/ICC** 1:50 - 1:200

## **Immunogen Information**

**Gene ID**Swiss Prot
115677
Q8IVI9

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 270-430 of human NOSTRIN (NP\_001034813.2).

## **Synonyms**

DaIP2; NOSTRIN

## **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**




Western blot analysis of various lysates using NOSTRIN Rabbit pAb (A15938) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunofluorescence analysis of U-2 OS cells using NOSTRIN Rabbit pAb (A15938) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.