

NeuN Rabbit pAb

Catalog No.: A15959

Basic Information

Observed MW

54kDa

Calculated MW

34kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the RNA-binding FOX protein family which is involved in the regulation of alternative splicing of pre-mRNA. The protein has an N-terminal proline-rich region, an RNA recognition motif (RRM) domain, and a C-terminal alanine-rich region. This gene produces the neuronal nuclei (NeuN) antigen that has been widely used as a marker for post-mitotic neurons. This gene has its highest expression in the central nervous system and plays a prominent role in neural tissue development and regulation of adult brain function. Mutations in this gene have been associated with numerous neurological disorders. Alternative splicing of this gene results in multiple transcript variants encoding distinct isoforms.

Recommended Dilutions

| | |
|--------------|----------------|
| WB | 1:500 - 1:2000 |
| IHC-P | 1:50 - 1:100 |

Immunogen Information

| | |
|----------------|-------------------|
| Gene ID | Swiss Prot |
| 146713 | A6NFN3 |

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of human NeuN (NP_001076044.1).

Synonyms

FOX3; NEUN; FOX-3; HRNBP3; NeuN

Contact

 | www.abclonal.com

Product Information


| | | |
|---------------|----------------|-----------------------|
| Source | Isotype | Purification |
| Rabbit | IgG | Affinity purification |

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from U-251MG cells, using NeuN Rabbit pAb (A15959) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 90s.



Immunohistochemistry analysis of NeuN in paraffin-embedded rat brain using NeuN Rabbit pAb (A15959) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of NeuN in paraffin-embedded mouse spinal cord using NeuN Rabbit pAb (A15959) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of NeuN in paraffin-embedded mouse brain using NeuN Rabbit pAb (A15959) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.