# ABclonal® www.abclonal.com

## **HSCB Rabbit pAb**

Catalog No.: A15961

## **Basic Information**

## **Observed MW**

27kDa

#### **Calculated MW**

27kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human

## **Background**

This gene encodes a DnaJ-type co-chaperone and member of the heat shock cognate B (HscB) family of proteins. The encoded protein plays a role in the synthesis of iron-sulfur clusters, protein cofactors that are involved in the redox reactions of mitochondrial electron transport and other processes. Cells in which this gene is knocked down exhibit reduced activity of iron-sulfur cluster-dependent enzymes including succinate dehydrogenase and aconitase. The encoded protein may stimulate the ATPase activity of the mitochondrial stress-70 protein. Alternative splicing results in multiple transcript variants.

## **Recommended Dilutions**

WB

1:500 - 1:2000

## **Immunogen Information**

Gene ID

**Swiss Prot** 

150274 Q8IWL3

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 30-235 of human HSCB (NP\_741999.3).

#### **Synonyms**

JAC1; HSC20; SIDBA5; DNAJC20; HSCB

## **Contact**

•

www.abclonal.com

## **Product Information**

**Source** Rabbit **Isotype** IgG Purification


Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using HSCB Rabbit pAb (A15961) at 1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.