

A16026

Leader in Biomolecular Solutions for Life Science



ZNRF3 Rabbit pAb

Catalog No.: A16026 **1 Publications**

Basic Information

Observed MW

101kDa

Calculated MW

101kDa

Category

Polyclonal Antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

Enables frizzled binding activity and ubiquitin-protein transferase activity. Involved in cellular protein metabolic process and negative regulation of Wnt signaling pathway. Is integral component of plasma membrane.

Recommended Dilutions

WB 1:1000 - 1:5000

IHC-P 1:50 - 1:200

Immunogen Information

Gene ID

84133

Swiss Prot

Q9ULT6

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 100-200 of human ZNRF3 (NP_001193927.1).

Synonyms

RNF203; BK747E2.3; ZNRF3

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.


Validation Data




Western blot analysis of lysates from Mouse testis, using ZNRF3 Rabbit pAb (A16026) at 1:3000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.



Immunohistochemistry analysis of ZNRF3 in paraffin-embedded human liver using ZNRF3 Rabbit pAb (A16026) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of ZNRF3 in paraffin-embedded mouse spleen using ZNRF3 Rabbit pAb (A16026) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of ZNRF3 in paraffin-embedded rat lung using ZNRF3 Rabbit pAb (A16026) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.