

δ-Catenin/p120 Catenin Rabbit pAb

Catalog No.: A1641

Basic Information

Observed MW

101kDa

Calculated MW

108kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the Armadillo protein family, which function in adhesion between cells and signal transduction. Multiple translation initiation codons and alternative splicing result in many different isoforms being translated. Not all of the full-length natures of the described transcript variants have been determined. Read-through transcription also exists between this gene and the neighboring upstream thioredoxin-related transmembrane protein 2 (TMX2) gene.

Recommended Dilutions

WB	1:500 - 1:2000
IHC-P	1:50 - 1:200

Immunogen Information

Gene ID	Swiss Prot
1500	O60716

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 850-968 of human δ-Catenin/p120 Catenin (NP_001078927.1).

Synonyms

CAS; p120; BCDS2; CTNND; P120CAS; P120CTN; p120(CAS); p120(CTN); δ-Catenin/p120 Catenin

Contact

 | www.abclonal.com


Product Information

Source	Isotype	Purification
Rabbit	IgG	Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from HT-29 cells, using δ -Catenin/p120 Catenin Rabbit pAb (A1641) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.




Immunohistochemistry analysis of δ -Catenin/p120 Catenin in paraffin-embedded human liver cancer using δ -Catenin/p120 Catenin Rabbit pAb (A1641) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of δ -Catenin/p120 Catenin in paraffin-embedded human prostate cancer using δ -Catenin/p120 Catenin Rabbit pAb (A1641) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of δ -Catenin/p120 Catenin in paraffin-embedded human urothelial carcinoma using δ -Catenin/p120 Catenin Rabbit pAb (A1641) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of δ -Catenin/p120 Catenin in paraffin-embedded mouse colon using δ -Catenin/p120 Catenin Rabbit pAb (A1641) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of δ -Catenin/p120 Catenin in paraffin-embedded rat uterus using δ -Catenin/p120 Catenin Rabbit pAb (A1641) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.