A17910

Leader in Biomolecular Solutions for Life Science



α -Smooth Muscle Actin (ACTA2) Rabbit mAb

Catalog No.: A17910 Recombinant

125 Publications

Basic Information

Observed MW 42kDa

Calculated MW 42kDa

Category SMab Recombinant Monoclonal Antibody

Applications WB,IF/ICC,ELISA

Cross-Reactivity Human,Mouse,Rat

CloneNo number ARC0298

Background

This gene encodes one of six different actin proteins. Actins are highly conserved proteins that are involved in cell motility, structure, integrity, and intercellular signaling. The encoded protein is a smooth muscle actin that is involved in vascular contractility and blood pressure homeostasis. Mutations in this gene cause a variety of vascular diseases, such as thoracic aortic disease, coronary artery disease, stroke, and Moyamoya disease, as well as multisystemic smooth muscle dysfunction syndrome.

Recommended Dilutions

WB	1:500 - 1:2000	Gene ID
17/100	1 50 1 200	59
IF/ICC	1:50 - 1:200	

Swiss Prot P62736

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human α -Smooth Muscle Actin (ACTA2) (P62736).

Synonyms

ACTSA; α-Smooth Muscle Actin (ACTA2)

€

Product Information

www.abclonal.com

lsotype IgG Purification Affinity purification


Storage

Source


Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates, using α -Smooth Muscle Actin (ACTA2) Rabbit mAb (A17910) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.



Confocal imaging of paraffin-embedded Human colon cancer tissue using α -Smooth Muscle Actin (ACTA2) Rabbit mAb (A17910, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x. Perform high pressure antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining.