A17917

Leader in Biomolecular Solutions for Life Science

Acetyl-Histone H3-K9/K14/K18/K23/K27 Rabbit pAb

ABclomal www.abclonal.com

Catalog No.: A17917 2 Publications

Basic Information

Observed MW 17kDa

Calculated MW 15kDa

Category Polyclonal Antibody

Applications WB, IF/ICC, ELISA

Cross-Reactivity Human, Mouse, Rat, Other (Wide Range Predicted)

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

Immunogen Information

WB	1:1000 - 1:5000	Gene ID
IF/ICC	1:50 - 1:200	8290/8350
		Immunogen

Swiss Prot Q16695/P68431

A synthetic acetylated peptide around K9 & K14 & K18 & K23 & K27 of human Histone H3 (NP 003520.1).

Synonyms

H3/A; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FA; H3C10; H3C11; H3C12; HIST1H3A; Acetyl-Histone H3-K9/K14/K18/K23/K27

Contact

Product Information

www.abclonal.com €

Isotype lgG

Purification Affinity purification


Storage

Source


Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from HeLa cells, using Acetyl-Histone H3-K9/K14/K18/K23/K27 Rabbit pAb (A17917) at 1:5000 dilution.HeLa cells were treated by TSA (1 uM) at 37°C for 18 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Immunofluorescence analysis of HeLa treated with TSA(1 uM,4 hours) and HeLa cells using Acetyl-Histone H3-K9/K14/K18/K23/K27 Rabbit pAb(A17917) at a dilution of 1:100 (40x lens). Secondary antibody:Cy3 Goat Anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.