Leader in Biomolecular Solutions for Life Science

pan-Myristyl-G Rabbit pAb

Catalog No.: A18599



Basic Information

Observed MW

12kDa/30kDa

Calculated MW

Category

Polyclonal Antibody

Applications

WB,ELISA

Cross-Reactivity

Rat

Background

Protein N-myristoylation is an important fatty acylation catalyzed by N-myristoyltransferases (NMTs), which are ubiquitous enzymes in eukaryotes. Specifically, attachment of a myristoyl group is vital for proteins participating in various biological functions, including signal transduction, cellular localization, and oncogenesis.

Recommended Dilutions

WB 1:100 - 1:500

Immunogen Information

Gene ID Swiss Prot

Immunogen

A synthetic peptide corresponding to a sequence containing myristylated G.

Synonyms

Contact

3

www.abclonal.com

Product Information

Source Rabbit **Isotype** IgG


PurificationAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from Rat liver, using pan-Myristyl-G Rabbit pAb (A18599) at 1:500

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.