

# Lamin B1 Rabbit pAb

Catalog No.: A1910 **40 Publications**

## Basic Information

### Observed MW

68kDa/45kDa

### Calculated MW

66kDa

### Category

Primary antibody

### Applications

ELISA, WB, IHC-P, IF/ICC, IP, CHIP

### Cross-Reactivity

Human, Mouse, Rat

## Background

This gene encodes one of the two B-type lamin proteins and is a component of the nuclear lamina. A duplication of this gene is associated with autosomal dominant adult-onset leukodystrophy (ADLD). Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

<b>WB</b>	1:500 - 1:1000
<b>IHC-P</b>	1:50 - 1:200
<b>IF/ICC</b>	1:50 - 1:200
<b>IP</b>	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
<b>ChIP</b>	5µg antibody for 10µg-15µg of Chromatin

## Contact

 | [www.abclonal.com](http://www.abclonal.com)

## Immunogen Information

<b>Gene ID</b>	<b>Swiss Prot</b>
4001	P20700

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 397-586 of human Lamin B1 (NP\_005564.1).

### Synonyms

LMN; ADLD; LMN2; LMNB; MCPH26; Lamin B1

## Product Information


<b>Source</b>	<b>Isotype</b>	<b>Purification</b>
Rabbit	IgG	Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


## Validation Data



Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg Lamin B1 antibody (A1910). Western blot was performed from the immunoprecipitate using Lamin B1 (A1910) at a dilution of 1:1000.




Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg Lamin B1 antibody (A1910). Western blot was performed from the immunoprecipitate using Lamin B1 antibody (A1910) at a dilution of 1:1000.




Western blot analysis of various lysates using Lamin B1 Rabbit pAb (A1910) at 1:1000 dilution. Jurkat cells were treated by Etoposide (25 µM) at 37°C for 5 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.


## Validation Data




Western blot analysis of various lysates using Lamin B1 Rabbit pAb (A1910) at 1:1000 dilution. L929 cells were treated by staurosporine (1  $\mu$ M) for 3 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25  $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Lamin B1 Rabbit pAb antibody (A1910) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Confocal immunofluorescence analysis of HeLa cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:200. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of PC-12 cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of U2OS cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

## Validation Data


---



Immunohistochemistry analysis of Lamin B1 in paraffin-embedded human breast cancer using Lamin B1 Rabbit pAb (A1910) at dilution of 1:150 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of Lamin B1 in paraffin-embedded mouse kidney using Lamin B1 Rabbit pAb (A1910) at dilution of 1:150 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of Lamin B1 in paraffin-embedded rat brain using Lamin B1 Rabbit pAb (A1910) at dilution of 1:150 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.