

A19292

Leader in Biomolecular Solutions for Life Science



## BCL2L14/BCLG Rabbit mAb

Catalog No.: A19292 **Recombinant**

### Basic Information

**Observed MW**

37kDa

**Calculated MW**

37kDa

**Category**

SMab Recombinant Monoclonal Antibody

**Applications**

WB,IHC-P,ELISA

**Cross-Reactivity**

Human,Mouse,Rat

**CloneNo number**

ARC2474

### Background

The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. Overexpression of this gene has been shown to induce apoptosis in cells. Three alternatively spliced transcript variants encoding two distinct isoforms have been reported for this gene.

### Recommended Dilutions

<b>WB</b>	1:500 - 1:1000
<b>IHC-P</b>	1:50 - 1:200

### Immunogen Information

**Gene ID**

79370

**Swiss Prot**

Q9BZR8

**Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human BCL2L14/BCLG (Q9BZR8).

**Synonyms**

BCLG; BCL2L14/BCLG

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**



Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.


## Validation Data




Western blot analysis of various lysates using BCL2L14/BCLG Rabbit mAb (A19292) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.



Immunohistochemistry analysis of BCL2L14/BCLG in paraffin-embedded mouse kidney tissue using BCL2L14/BCLG Rabbit mAb (A19292) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of BCL2L14/BCLG in paraffin-embedded human colon tissue using BCL2L14/BCLG Rabbit mAb (A19292) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of BCL2L14/BCLG in paraffin-embedded mouse testis tissue using BCL2L14/BCLG Rabbit mAb (A19292) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.

## Validation Data


---



Immunohistochemistry analysis of BCL2L14/BCLG in paraffin-embedded rat testis tissue using BCL2L14/BCLG Rabbit mAb (A19292) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of BCL2L14/BCLG in paraffin-embedded Human lung adenocarcinoma tissue using BCL2L14/BCLG Rabbit mAb (A19292) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of BCL2L14/BCLG in paraffin-embedded rat kidney tissue using BCL2L14/BCLG Rabbit mAb (A19292) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.