

A19754

Leader in Biomolecular Solutions for Life Science



alpha Sarcoglycan (SGCA) Rabbit mAb

Catalog No.: A19754 **Recombinant**

Basic Information

Observed MW

48kDa

Calculated MW

43kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity

Mouse,Rat

CloneNo number

ARC2280

Background

This gene encodes a component of the dystrophin-glycoprotein complex (DGC), which is critical to the stability of muscle fiber membranes and to the linking of the actin cytoskeleton to the extracellular matrix. Its expression is thought to be restricted to striated muscle. Mutations in this gene result in type 2D autosomal recessive limb-girdle muscular dystrophy. Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:100 - 1:500

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

Immunogen Information

Gene ID

6442

Swiss Prot

Q16586

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 288-387 of human alpha Sarcoglycan (SGCA) (SGCA) (Q16586).

Synonyms

ADL; DAG2; 50DAG; DMDA2; LGMD2D; LGMDR3; SCARMD1; adhalin; alpha Sarcoglycan (SGCA)

Contact

 www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using alpha Sarcoglycan (SGCA) (SGCA) Rabbit mAb (A19754) at 1:500 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.




Immunohistochemistry analysis of paraffin-embedded mouse heart using alpha Sarcoglycan (SGCA) (SGCA) Rabbit mAb (A19754) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat heart using alpha Sarcoglycan (SGCA) (SGCA) Rabbit mAb (A19754) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Confocal imaging of paraffin-embedded Mouse heart using alpha Sarcoglycan (SGCA) Rabbit mAb (A19754, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x. Perform high pressure antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining.



Confocal imaging of paraffin-embedded Rat heart using alpha Sarcoglycan (SGCA) Rabbit mAb (A19754, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used

Validation Data

for nuclear staining (Blue). Objective:
40x. Perform high pressure antigen
retrieval with 0.01 M citrate buffer
(pH 6.0) prior to IF staining.