# OsMADS1 Rabbit pAb

Catalog No.: A20328 2 Publications



### **Basic Information**

**Observed MW** Refer to figures

Calculated MW 30kDa

**Category** Polyclonal Antibody

Applications WB,ELISA

Cross-Reactivity Oryza sativa

# Background

Probable transcription factor involved in the development of floral organs. Required for the formation of inner floral organs (lodicules, stamens and carpels, or whorls 2, 3 and 4 and the lemma and palea (whorl 1, which are grass floral organs analogous to sepals. May be involved in the control of flowering time. Seems to act as transcriptional activator. May act upstream of the auxin-responsive protein GH3.8.

# Immunogen Information

| WB | 1:500 - 1:2000 | <b>Gene ID</b><br>4332059   | Swiss Prot<br>Q10PZ9 |
|----|----------------|---|----------------------|
|    |                | <b>Immunogen</b><br>Recombinant fusion protein containing a sequence corresponding to amino acids |                      |

Recombinant fusion protein containing a sequence corresponding to amino acids 150-257 of Oryza sativa OsMADS1. (Q10PZ9).

#### Synonyms

Lhs1; MADS1; OsMADS1; OsJ\_09913

### **Product Information**

www.abclonal.com

**Recommended Dilutions** 

**Isotype** IgG **Purification** Affinity purification


#### Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of extracts of various tissues from the japonica rice (Oryza sativa L.) variety Zhonghua 11, using OsMADS1 Rabbit pAb (A20328) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 120s.