# ABclonal® www.abclonal.com

## OsGSK2 Rabbit pAb

Catalog No.: A20367

## **Basic Information**

#### **Observed MW**

45kDa

#### **Calculated MW**

46kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,ELISA

## **Cross-Reactivity**

Oryza sativa

## **Background**

Serine-threonine kinase that acts as a negative regulator of brassinosteroid (BR) signaling. Phosphorylates DLT and BZR1, two positive regulators that mediates several BR responses. Phosphorylation of DLT and BZR1 inhibits their activities in BR signaling. Phosphorylates OFP8, a positive regulator of BR responses. Phosphorylated OFP8 shuttles from the nucleus to the cytoplasm where it is degraded by the proteasome. Phosphorylates the E3 ubiquitin-protein ligase PUB24, a negative regulator of BR signaling, which targets BZR1 and promotes its degradation via the 26S proteasome. Phosphorylation of PUB24 increases its stability. Phosphorylates the AP2-ERF transcription factor SMOS1, a positive regulator of BR signaling, which cooperatively functions in a transactivating complex with BZR1 to enhance the transcription of BR biosynthetic genes. Phosphorylation of SMOS1 leads to its degradation by an unknown mechanism.

## **Recommended Dilutions**

**WB** 

1:500 - 1:2000

## **Immunogen Information**

Gene ID

Swiss Prot Q60EZ2

4338079

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1-100 of Oryza sativa OsGSK2. (Q60EZ2).

## **Synonyms**

GSK2; SK22; OsGSK2; OsSK22; OJ1430\_B02.4

## **Contact**

•

www.abclonal.com


## **Product Information**

**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.



Western blot analysis of extracts of various tissues from the japonica rice (Oryza sativa L.) variety Zhonghua 11, using OsGSK2 Rabbit pAb (A20367) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.