

A2353

Leader in Biomolecular Solutions for Life Science



# CD86 Rabbit pAb

Catalog No.: A2353 **3 Publications**

## Basic Information

### Observed MW

70kDa

### Calculated MW

38kDa

### Category

Polyclonal Antibody

### Applications

WB,IHC-P,ELISA

### Cross-Reactivity

Mouse

## Background

This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript variants encoding different isoforms.

## Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:100

## Immunogen Information

### Gene ID

942

### Swiss Prot

P42081

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 24-247 of human CD86 (NP\_787058.4).

### Synonyms

B70; B7-2; B7.2; LAB72; CD28LG2; CD86

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of various lysates using CD86 Rabbit pAb (A2353) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 300s.



Immunohistochemistry analysis of CD86 in paraffin-embedded mouse spleen using CD86 Rabbit pAb (A2353) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.