

iNOS Rabbit pAb

Catalog No.: A3200 **8 Publications**

Basic Information

Observed MW

131kDa

Calculated MW

131kDa

Category

Primary antibody

Applications

ELISA, WB

Cross-Reactivity

Human, Mouse

Background

Nitric oxide is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission and antimicrobial and antitumoral activities. This gene encodes a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. Three related pseudogenes are located within the Smith-Magenis syndrome region on chromosome 17.

Recommended Dilutions

WB 1:500 - 1:1000

Immunogen Information

Gene ID

4843

Swiss Prot

P35228

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-170 of human iNOS (NP_000616.3).

Synonyms

NOS; INOS; NOS2A; HEP-NOS; iNOS

Contact

 | www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of RAW264.7 cells, using iNOS antibody (A3200) at 1:1000 dilution. Raw264.7 cells were treated by LPS (1 µg/ml) at 37°C for over night.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.



Western blot analysis of extracts of RAW264.7 cells, using iNOS antibody (A3200) at 1:1000 dilution. Raw264.7 cells were treated by LPS (1 µg/ml) at 37°C for 8 hours.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.