# ABclonal www.abclonal.com

# **GULP1 Rabbit pAb**

Catalog No.: A3432

# **Basic Information**

#### **Observed MW**

34kDa

# **Calculated MW**

34kDa

#### Category

Primary antibody

# **Applications**

ELISA, WB, IHC-P

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene is an adapter protein necessary for the engulfment of apoptotic cells by phagocytes. Several transcript variants, some protein coding and some thought not to be protein coding, have been found for this gene.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IHC-P** 1:50 - 1:200

# **Immunogen Information**

**Gene ID**51454

Swiss Prot
Q9UBP9

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-130 of human GULP1 (NP\_057399.1).

#### **Synonyms**

CED6; GULP; CED-6; GULP1

# **Contact**

www.abclonal.com


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.




Western blot analysis of various lysates using GULP1 Rabbit pAb (A3432) at 1:3000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).


Exposure time: 90s.



Immunohistochemistry analysis of GULP1 in paraffin-embedded Mouse spinal cord using GULP1 Rabbit pAb (A3432) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of GULP1 in paraffin-embedded Rat ovary using GULP1 Rabbit pAb (A3432) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of GULP1 in paraffin-embedded Human colon carcinoma using GULP1 Rabbit pAb (A3432) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.