Leader in Biomolecular Solutions for Life Science

# ATP6V1B2 Rabbit pAb

Catalog No.: A3754



### **Basic Information**

### **Observed MW**

57kDa

#### **Calculated MW**

57kDa

#### Category

Mouse Monoclonal Antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A, three B, and two G subunits, as well as a C, D, E, F, and H subunit. The V1 domain contains the ATP catalytic site. The protein encoded by this gene is one of two V1 domain B subunit isoforms and is the only B isoform highly expressed in osteoclasts.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# Immunogen Information

**Gene ID** 

526

Swiss Prot P21281

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 262-511 of human ATP6V1B2 (NP 001684.2).

### **Synonyms**

DOOD; HO57; VATB; VPP3; Vma2; ZLS2; ATP6B2; ATP6B1B2; ATP6V1B2

### **Contact**

•

www.abclonal.com

## **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using ATP6V1B2 Antibody (A3754) at 1:1000

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.