# Troponin I2 (TNNI2) Rabbit pAb

Catalog No.: A4162



# **Basic Information**

**Observed MW** 21kDa

Calculated MW 21kDa

**Category** Primary antibody

Applications WB

Cross-Reactivity Human

### Background

This gene encodes a fast-twitch skeletal muscle protein, a member of the troponin I gene family, and a component of the troponin complex including troponin T, troponin C and troponin I subunits. The troponin complex, along with tropomyosin, is responsible for the calcium-dependent regulation of striated muscle contraction. Mouse studies show that this component is also present in vascular smooth muscle and may play a role in regulation of smooth muscle function. In addition to muscle tissues, this protein is found in corneal epithelium, cartilage where it is an inhibitor of angiogenesis to inhibit tumor growth and metastasis, and mammary gland where it functions as a co-activator of estrogen receptor-related receptor alpha. This protein also suppresses tumor growth in human ovarian carcinoma. Mutations in this gene cause myopathy and distal arthrogryposis type 2B. Alternatively spliced transcript variants have been found for this gene.

### **Recommended Dilutions**

1:500 - 1:1000

# Immunogen Information

WB

**Gene ID** 7136 Swiss Prot P48788

#### Immunogen

A synthetic Peptide of human Troponin I2 (Troponin I2 (TNNI2))

Isotype

lgG

Synonyms

DA2B; FSSV; DA2B1; fsTnI; AMCD2B; Troponin I2 (TNNI2)

# **Product Information**

 www.abclonal.com


Purification Affinity purification

Storage

Source

Rabbit

Store at 4°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,pH7.3.



Western blot analysis of extracts of human skeletal muscle, using Troponin I2 (Troponin I2 (TNNI2)) antibody (A4162).

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.