# ABclonal www.abclonal.com

## **ABCG1 Rabbit pAb**

Catalog No.: A4328 6 Publications

## **Basic Information**

#### **Observed MW**

75kDa

## **Calculated MW**

76kDa

## Category

Primary antibody

## **Applications**

ELISA, WB, IHC-P, IF/ICC

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. It is involved in macrophage cholesterol and phospholipids transport, and may regulate cellular lipid homeostasis in other cell types. Six alternative splice variants have been identified.

## **Recommended Dilutions**

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:100

## **Immunogen Information**

Gene ID	<b>Swiss Prot</b>
9619	P45844

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 6-120 of human ABCG1 (NP\_004906.3).

#### **Synonyms**

ABC8; WHITE1; ABCG1

## **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates, using ABCG1 Rabbit pAb (A4328) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20s.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma using ABCG1 Rabbit pAb (A4328) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of Rat lung using ABCG1 antibody (A4328) at dilution of 1:100. Blue: DAPI for nuclear staining.