# ABclonal www.abclonal.com

## **NUP93 Rabbit pAb**

Catalog No.: A4333 1 Publications

## **Basic Information**

#### **Observed MW**

93kDa

#### **Calculated MW**

93kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

The nuclear pore complex is a massive structure that extends across the nuclear envelope, forming a gateway that regulates the flow of macromolecules between the nucleus and the cytoplasm. Nucleoporins are the main components of the nuclear pore complex in eukaryotic cells. This gene encodes a nucleoporin protein that localizes both to the basket of the pore and to the nuclear entry of the central gated channel of the pore. The encoded protein is a target of caspase cysteine proteases that play a central role in programmed cell death by apoptosis. Alternative splicing results in multiple transcript variants encoding different isoforms.

## **Recommended Dilutions**

WB

1:500 - 1:2000

## Immunogen Information

Gene ID 9688 **Swiss Prot** 

Q8N1F7

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 670-819 of human NUP93 (NP\_055484.3).

#### **Synonyms**

NIC96; NUP93

## **Contact**

•

www.abclonal.com

## **Product Information**

**Source** Rabbit Isotype

IgG

**Purification** 


Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using NUP93 Rabbit pAb (A4333) at 1:3000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.