Leader in Biomolecular Solutions for Life Science

# **NUP93 Rabbit pAb**

Catalog No.: A4333 1 Publications



### **Basic Information**

### **Observed MW**

93kDa

#### **Calculated MW**

93kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The nuclear pore complex is a massive structure that extends across the nuclear envelope, forming a gateway that regulates the flow of macromolecules between the nucleus and the cytoplasm. Nucleoporins are the main components of the nuclear pore complex in eukaryotic cells. This gene encodes a nucleoporin protein that localizes both to the basket of the pore and to the nuclear entry of the central gated channel of the pore. The encoded protein is a target of caspase cysteine proteases that play a central role in programmed cell death by apoptosis. Alternative splicing results in multiple transcript variants encoding different isoforms.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

Gene ID

**Swiss Prot** 

Q8N1F7

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 670-819 of human NUP93 (NP 055484.3).

# Synonyms

NIC96; NUP93

### **Contact**

•

www.abclonal.com

## **Product Information**

**Source** Rabbit **Isotype** IgG Purification


Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using NUP93 Rabbit pAb (A4333) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.