

MIER2 Rabbit pAb

Catalog No.: A4788

Basic Information

Observed MW

55kDa

Calculated MW

60kDa

Category

Primary antibody

Applications

WB,FC

Cross-Reactivity

Human

Background

Enables histone deacetylase binding activity. Contributes to histone deacetylase activity. Involved in histone deacetylation. Located in cytoplasm and nucleus. Part of protein-containing complex.

Recommended Dilutions

WB	1:500 - 1:1000
FC	1:20 - 1:50

Immunogen Information

Gene ID	Swiss Prot
54531	Q8N344

Immunogen

A synthetic Peptide of human MIER2

Synonyms

Mi-er2; KIAA1193; MIER2

Contact

 | www.abclonal.com


Product Information

Source	Isotype	Purification
Rabbit	IgG	Affinity purification

Storage

Store at 4°C. Avoid freeze / thaw cycles.
Buffer: PBS with 0.02% sodium azide,pH7.3.

Validation Data



Western blot analysis of extracts of NCI-H460 cells, using MIER2 antibody (A4788).
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.