

# ATRIP Rabbit pAb

Catalog No.: A5041

## Basic Information

### Observed MW

100kDa/113kDa

### Calculated MW

86kDa

### Category

Primary antibody

### Applications

ELISA, WB, IF/ICC, IP

### Cross-Reactivity

Human

## Background

This gene encodes an essential component of the DNA damage checkpoint. The encoded protein binds to single-stranded DNA coated with replication protein A. The protein also interacts with the ataxia telangiectasia and Rad3 related protein kinase, resulting in its accumulation at intranuclear foci induced by DNA damage. Multiple transcript variants encoding different isoforms have been found for this gene.

## Recommended Dilutions

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:200

**IP** 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells

## Immunogen Information

### Gene ID

84126

### Swiss Prot

Q8WXE1

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human ATRIP (NP\_569055.1).

### Synonyms

ATRIP

## Contact

 | [www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

# Validation Data



Immunoprecipitation analysis of 200 µg extracts of HeLa cells using 1 µg ATRIP antibody (A5041). Western blot was performed from the immunoprecipitate using ATRIP antibody (A5041) at a dilution of 1:1000.



Western blot analysis of various lysates using ATRIP Rabbit pAb (A5041) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.