Leader in Biomolecular Solutions for Life Science

# **SULT2A1** Rabbit pAb

Catalog No.: A5489



### **Basic Information**

### **Observed MW**

34kDa

#### **Calculated MW**

34kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse

### **Background**

This gene encodes a member of the sulfotransferase family. Sulfotransferases aid in the metabolism of drugs and endogenous compounds by converting these substances into more hydrophilic water-soluble sulfate conjugates that can be easily excreted. This protein catalyzes the sulfation of steroids and bile acids in the liver and adrenal glands, and may have a role in the inherited adrenal androgen excess in women with polycystic ovary syndrome.

### **Recommended Dilutions**

WB

1:500 - 1:2000

### **Immunogen Information**

Gene ID 6822 Swiss Prot Q06520

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-145 of human SULT2A1 ( $NP_003158.2$ ).

### **Synonyms**

HST; ST2; STD; hSTa; DHEAS; ST2A1; ST2A3; DHEA-ST; SULT2A3; DHEA-ST8; SULT2A1

### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of extracts of various cell lines, using SULT2A1 antibody (A5489) at 1:2000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.