# ABclonal www.abclonal.com

# KITLG Rabbit pAb

Catalog No.: A5672 1 Publications

## **Basic Information**

#### **Observed MW**

31kDa

#### **Calculated MW**

31kDa

#### Category

Polyclonal Antibody

## **Applications**

WB,IF/ICC,ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes the ligand of the tyrosine-kinase receptor encoded by the KIT locus. This ligand is a pleiotropic factor that acts in utero in germ cell and neural cell development, and hematopoiesis, all believed to reflect a role in cell migration. In adults, it functions pleiotropically, while mostly noted for its continued requirement in hematopoiesis. Two transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

WB 1:500 - 1:1000

IF/ICC 1:50 - 1:200

# **Immunogen Information**

**Gene ID Swiss Prot** 4254 P21583

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 120-190 of human KITLG (NP\_000890.1).

### **Synonyms**

SF; MGF; SCF; SLF; DCUA; FPH2; FPHH; KL-1; Kitl; WS2F; SHEP7; DFNA69; KITLG

## **Contact**

0 www.abclonal.com


## **Product Information**

Source **Isotype Purification** Rabbit Affinity purification IgG

## Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of extracts of various cell lines, using KITLG antibody (A5672) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Immunofluorescence analysis of NIH/3T3 cells using KITLG Rabbit pAb (A5672) at dilution of 1:20 (40x lens). Blue: DAPI for nuclear staining.