Leader in Biomolecular Solutions for Life Science

# **ACTR1A Rabbit pAb**

Catalog No.: A6515



### **Basic Information**

### **Observed MW**

43kDa

### **Calculated MW**

43kDa

### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

### **Background**

This gene encodes a 42.6 kD subunit of dynactin, a macromolecular complex consisting of 10-11 subunits ranging in size from 22 to 150 kD. Dynactin binds to both microtubules and cytoplasmic dynein. It is involved in a diverse array of cellular functions, including ER-to-Golgi transport, the centripetal movement of lysosomes and endosomes, spindle formation, chromosome movement, nuclear positioning, and axonogenesis. This subunit is present in 8-13 copies per dynactin molecule, and is the most abundant molecule in the dynactin complex. It is an actin-related protein, and is approximately 60% identical at the amino acid level to conventional actin.

### **Recommended Dilutions**

WB

1:500 - 1:2000

### **Immunogen Information**

**Gene ID** 10121

**Swiss Prot** 

P61163

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 187-376 of human ACTR1A (NP\_005727.1).

### **Synonyms**

ARP1; Arp1A; CTRN1; ACTR1A

### **Contact**

•

www.abclonal.com

### **Product Information**

**Source** Rabbit **Isotype** IgG

Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of extracts of various cell lines, using ACTR1A antibody (A6515) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.