

A7255

Leader in Biomolecular Solutions for Life Science



# Acetyl-Histone H3-K9 Rabbit pAb

Catalog No.: A7255

65 Publications

## Basic Information

### Observed MW

17kDa

### Calculated MW

16kDa

### Category

Polyclonal Antibody

### Applications

WB, IHC-P, IF/ICC, IP, ChIP, ChIP-seq, ELISA

### Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

## Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200
IP	0.5ug-4ug antibody for 200ug-400ug extracts of whole cells
ChIP	5µg antibody for 5µg-10µg of Chromatin
ChIP-seq	1:20 - 1:50

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Immunogen Information

### Gene ID

8290/8350

### Swiss Prot

Q16695/P68431

### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Histone H3 (NP\_003520.1).

### Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; Acetyl-Histone H3-K9

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.


## Validation Data



Western blot analysis of lysates from HeLa cells, using Acetyl-Histone H3-K9 Rabbit pAb (A7255) at 1:1000 dilution. HeLa cells were treated by TSA (1  $\mu$ M) at 37°C for 18 hours.  
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25  $\mu$ g per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 10s.




Western blot analysis of lysates from NIH/3T3 cells, using Acetyl-Histone H3-K9 Rabbit pAb (A7255) at 1:1000 dilution. NIH/3T3 cells were treated by TSA (1  $\mu$ M) at 37°C for 18 hours.  
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25  $\mu$ g per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 10s.




Western blot analysis of lysates from C6 cells, using Acetyl-Histone H3-K9 Rabbit pAb (A7255) at 1:1000 dilution. C6 cells were treated by TSA (1  $\mu$ M) at 37°C for 18 hours.  
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25  $\mu$ g per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 10s.


## Validation Data




Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded human mammary cancer using Acetyl-Histone H3-K9 Rabbit pAb (A7255) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded human colon using Acetyl-Histone H3-K9 Rabbit pAb (A7255) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded rat ovary using Acetyl-Histone H3-K9 Rabbit pAb (A7255) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded mouse brain using Acetyl-Histone H3-K9 Rabbit pAb (A7255) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of HeLa cells using Acetyl-Histone H3-K9 Rabbit pAb (A7255) at dilution of 1:100 (40x lens). HeLa cells were treated by TSA (1  $\mu$ M) at 37°C for 18 hours (left). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using Acetyl-Histone H3-K9 Rabbit pAb (A7255) at dilution of 1:100 (40x lens). NIH/3T3 cells were treated by TSA (1  $\mu$ M) at 37°C for 18 hours (left). Blue: DAPI for nuclear staining.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Acetyl-Histone H3-K9 antibody (A7255) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.