Leader in Biomolecular Solutions for Life Science

# **ABCG5 Rabbit pAb**

Catalog No.: A8589 1 Publications



### **Basic Information**

### **Observed MW**

80kDa

### **Calculated MW**

73kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human

### **Background**

The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. The protein encoded by this gene functions as a half-transporter to limit intestinal absorption and promote biliary excretion of sterols. It is expressed in a tissue-specific manner in the liver, colon, and intestine. This gene is tandemly arrayed on chromosome 2, in a head-to-head orientation with family member ABCG8. Mutations in this gene may contribute to sterol accumulation and atheroschlerosis, and have been observed in patients with sitosterolemia.

### **Recommended Dilutions**

WB

1:500 - 1:2000

### **Immunogen Information**

**Gene ID** 

Swiss Prot 09H222

64240

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 350-450 of human ABCG5 (NP 071881.1).

### **Synonyms**

STSL; STSL2; ABCG5

### **Contact**

•

www.abclonal.com

### **Product Information**

**Source** Rabbit **Isotype** IgG Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of extracts of HL-60 cells, using ABCG5 antibody (A8589) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.