Leader in Biomolecular Solutions for Life Science

COX7A1 Rabbit pAb

Catalog No.: A8707



Basic Information

Observed MW

10kDa

Calculated MW

9kDa

Category

Polyclonal Antibody

Applications

WB

Cross-Reactivity

Human, Mouse

Background

Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes polypeptide 1 (muscle isoform) of subunit VIIa and the polypeptide 1 is present only in muscle tissues. Other polypeptides of subunit VIIa are present in both muscle and nonmuscle tissues, and are encoded by different genes.

Recommended Dilutions

WB

1:500 - 1:1000

Immunogen Information

Gene ID

Swiss Prot P24310

1346

ImmunogenA synthetic Peptide of human COX7A1

Synonyms

COX7A; COX7AH; COX7AM; COX7A1

Contact

•

www.abclonal.com


Product Information

Source Rabbit **Isotype** IgG **Purification**Affinity purification

Storage

Store at 4°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using COX7A1 antibody (A8707). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.