

# Mouse anti TAP-Tag mAb

Catalog No.: AE021

## Basic Information

---

### Observed MW

Refer to Figures

### Calculated MW

### Category

Tag antibody

### Applications

WB,IP

### Cross-Reactivity

### CloneNo number

AMC0510

## Background

---

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

## Recommended Dilutions

---

<b>WB</b>	1:5000 - 1:10000
<b>IP</b>	1:100 - 1:200

## Immunogen Information

---

<b>Gene ID</b>	<b>Swiss Prot</b>
----------------	-------------------

### Immunogen

A synthetic peptide of TAP-Tag

### Synonyms

TAP;TAP tag;TAP-tag

## Contact

---

 | [www.abclonal.com](http://www.abclonal.com)

## Product Information

---


<b>Source</b>	<b>Isotype</b>	<b>Purification</b>
Mouse	IgG1	Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.  
Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## Validation Data

---



Western blot analysis of recombinant TAP protein using mouse anti TAP-Tag mAb (AE021) at dilution of 1:5000.

Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.