Leader in Biomolecular Solutions for Life Science

# Phospho-MAX-S11 Rabbit pAb

Catalog No.: AP0072



### **Basic Information**

### **Observed MW**

21kDa

#### **Calculated MW**

18kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human

# **Background**

The protein encoded by this gene is a member of the basic helix-loop-helix leucine zipper (bHLHZ) family of transcription factors. It is able to form homodimers and heterodimers with other family members, which include Mad, Mxi1 and Myc. Myc is an oncoprotein implicated in cell proliferation, differentiation and apoptosis. The homodimers and heterodimers compete for a common DNA target site (the E box) and rearrangement among these dimer forms provides a complex system of transcriptional regulation. Mutations of this gene have been reported to be associated with hereditary pheochromocytoma. A pseudogene of this gene is located on the long arm of chromosome 7. Alternative splicing results in multiple transcript variants.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

Gene ID

Swiss Prot P61244

Immunogen

A synthetic phosphorylated peptide around S11 of human MAX (NP\_002373.3).

### **Synonyms**

bHLHd4; Phospho-MAX-S11

### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from Jurkat cells, using Phospho-MAX-S11 Rabbit pAb (AP0072) at 1:1000 dilution. Jurkat cells were treated by serum-starvation overnight.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% BSA.