AP0121

Leader in Biomolecular Solutions for Life Science

Phospho-MEK2-T394 Rabbit pAb

Catalog No.: AP0121 1 Publications



Basic Information

Observed MW 44kDa

Calculated MW 44kDa

Category Polyclonal Antibody

Applications WB,IF/ICC

Cross-Reactivity Human

Background

The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinases. Mutations in this gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, cognitive disability, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene.

Recommended Dilutions

Immunogen Information

WB	1:500 - 1:2000
IF/ICC	1:100 - 1:200

Gene ID	
5605	

Swiss Prot P36507

Immunogen

A phospho specific peptide corresponding to residues surrounding T394 of human MEK2

Synonyms

CFC4; MEK2; MKK2; MAPKK2; PRKMK2; Phospho-MEK2-T394

€

Product Information

www.abclonal.com

lsotype IgG Purification Affinity purification


Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from HepG2 cells using Phospho-MEK2-T394 Rabbit pAb (AP0121). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA.