

AP0182

Leader in Biomolecular Solutions for Life Science



Phospho-LCK-Y394 Rabbit pAb

Catalog No.: AP0182 **3 Publications**

Basic Information

Observed MW

Calculated MW
58kDa

Category
Polyclonal Antibody

Applications
WB,IF/ICC,ELISA

Cross-Reactivity
Human,Mouse

Background

This gene is a member of the Src family of protein tyrosine kinases (PTKs). The encoded protein is a key signaling molecule in the selection and maturation of developing T-cells. It contains N-terminal sites for myristylation and palmitoylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to the plasma membrane and pericentrosomal vesicles, and binds to cell surface receptors, including CD4 and CD8, and other signaling molecules. Multiple alternatively spliced variants encoding different isoforms have been described.

Recommended Dilutions

WB	1:500 - 1:1000
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID	Swiss Prot
3932	P06239

Immunogen
A synthetic phosphorylated peptide around Y394 of human LCK (NP_005347.3).

Synonyms
LSK; YT16; IMD22; p56lck; pp58lck; Phospho-LCK-Y394

Contact


 | www.abclonal.com

Product Information

Source	Isotype	Purification
Rabbit	IgG	Affinity purification



Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from NIH/3T3 cells, using Phospho-LCK-Y394 Rabbit pAb (AP0182) at 1:1000 dilution. NIH/3T3 cells were treated by Pervanadate (1 mM) at 37°C for 30 minutes after serum-starvation overnight.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 180s.

Western blot analysis of lysates from Jurkat cells, using Phospho-LCK-Y394 Rabbit pAb (AP0182) at 1:1000 dilution. Jurkat cells were treated by CIP (20uL/400ul) at 37°C for 1 hour.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Enhanced Kit (RM00021).
 Exposure time: 180s.



Immunofluorescence analysis of Raji pervanadate and Raji cells using Phospho-LCK-Y394 Rabbit pAb (AP0182) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.