Leader in Biomolecular Solutions for Life Science

# Phospho-ABL1-Y393/ABL2-Y429 Rabbit pAb



Catalog No.: AP0282

# **Basic Information**

## **Observed MW**

210kDa

#### **Calculated MW**

122kDa/124kDa/60kDa/114-128kDa

### Category

Polyclonal Antibody

#### **Applications**

WB,IF/ICC

#### **Cross-Reactivity**

Human, Mouse

# **Background**

This gene is a protooncogene that encodes a protein tyrosine kinase involved in a variety of cellular processes, including cell division, adhesion, differentiation, and response to stress. The activity of the protein is negatively regulated by its SH3 domain, whereby deletion of the region encoding this domain results in an oncogene. The ubiquitously expressed protein has DNA-binding activity that is regulated by CDC2-mediated phosphorylation, suggesting a cell cycle function. This gene has been found fused to a variety of translocation partner genes in various leukemias, most notably the t(9;22) translocation that results in a fusion with the 5' end of the breakpoint cluster region gene (BCR; MIM:151410). Alternative splicing of this gene results in two transcript variants, which contain alternative first exons that are spliced to the remaining common exons. [provided by RefSeq, Aug 2014]

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

IF/ICC 1:100 - 1:200

# **Immunogen Information**

**Gene ID Swiss Prot**25/27
P00519/P42684

#### **Immunogen**

A phospho specific peptide corresponding to residues surrounding Y393/429 of human ABL1/2

## **Synonyms**

ABL1/ABL2; Phospho-ABL1-Y393/ABL2-Y429

### **Contact**

www.abclonal.com

# **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts from HL60 cells using phospho-ABL1/2-Y393/429 antibody (AP0282) and the same antibody preincubated with blocking peptide.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% BSA.