

AP0307

Leader in Biomolecular Solutions for Life Science



Phospho-Androgen Receptor-S650 Rabbit pAb

Catalog No.: AP0307

Basic Information

Observed MW

Calculated MW

99kDa

Category

Polyclonal Antibody

Applications

Cross-Reactivity

Human

Background

The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expansion of the polyglutamine tract from the normal 9-34 repeats to the pathogenic 38-62 repeats causes spinal bulbar muscular atrophy (SBMA, also known as Kennedy's disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Alternative splicing results in multiple transcript variants encoding different isoforms.

Recommended Dilutions

Immunogen Information

Gene ID

367

Swiss Prot

P10275

Immunogen

A phospho specific peptide corresponding to residues surrounding S650 of human AR

Synonyms

KD; AIS; AR8; TFM; DHTR; SBMA; HYSP1; NR3C4; SMAX1; HUMARA; Phospho-Androgen Receptor-S650

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

