Leader in Biomolecular Solutions for Life Science

# ABclonal www.abclonal.com

# Phospho-CDKN1A/p21CIP1-T145 Rabbit pAb

Catalog No.: AP0327

#### **Basic Information**

**Observed MW** 

Calculated MW 18kDa

**Category**Polyclonal Antibody

**Applications** 

Cross-Reactivity
Human

# **Background**

This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or -cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Multiple alternatively spliced variants have been found for this gene.

# **Recommended Dilutions**

# **Immunogen Information**

Gene ID

Swiss Prot P38936

**Immunogen** 

A phospho specific peptide corresponding to residues surrounding T145 of human CDKN1A/p21CIP1

**Synonyms** 

P21; CIP1; SDI1; WAF1; CAP20; CDKN1; MDA-6; p21CIP1; Phospho-CDKN1A/p21CIP1-T145

#### **Contact**

€

www.abclonal.com

## **Product Information**

**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.