Leader in Biomolecular Solutions for Life Science

Phospho-CDKN1B/p27KIP1-T187 Rabbit pAb



Catalog No.: AP0328

Basic Information

Observed MW

26kDa

Calculated MW

22kDa

Category

Polyclonal Antibody

Applications

WB

Cross-Reactivity

Human

Background

This gene encodes a cyclin-dependent kinase inhibitor, which shares a limited similarity with CDK inhibitor CDKN1A/p21. The encoded protein binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state. Mutations in this gene are associated with multiple endocrine neoplasia type IV (MEN4).

Recommended Dilutions

WB

1:500 - 1:1000

Immunogen Information

Gene ID

Swiss Prot P46527

Immunogen

A phospho specific peptide corresponding to residues surrounding T187 of human CDKN1B/p27KIP1

Synonyms

KIP1; MEN4; CDKN4; MEN1B; P27KIP1; Phospho-CDKN1B/p27KIP1-T187

Contact

•

www.abclonal.com

Product Information


Source Rabbit **Isotype** IgG **Purification**Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



 $We stern \ blot \ analysis \ of \ lysates \ from \ He La \ cells, \ using \ Phospho-CDKN1B/p27KIP1-T187 \ Rabbit \ pAb$

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% BSA.