Leader in Biomolecular Solutions for Life Science

Phospho-MARCKS-S162 Rabbit pAb

Catalog No.: AP0403



Basic Information

Observed MW

75kDa

Calculated MW

32kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

The protein encoded by this gene is a substrate for protein kinase C. It is localized to the plasma membrane and is an actin filament crosslinking protein. Phosphorylation by protein kinase C or binding to calcium-calmodulin inhibits its association with actin and with the plasma membrane, leading to its presence in the cytoplasm. The protein is thought to be involved in cell motility, phagocytosis, membrane trafficking and mitogenesis.

Recommended Dilutions

WB

1:100 - 1:500

Immunogen Information

Gene ID 4082 Swiss Prot P29966

Immunogen

A phospho specific peptide corresponding to residues surrounding S162 of human MARCKS

Synonyms

MACS; 80K-L; PKCSL; PRKCSL; Phospho-MARCKS-S162

Contact

•

www.abclonal.com

Product Information


Source Rabbit **Isotype** IgG **Purification**Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates, using Phospho-MARCKS-S162 antibody (AP0403) at 1:400 dilution.293T cells were treated by PMA/TPA (200 nM) at 37° C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.