Leader in Biomolecular Solutions for Life Science

# Phospho-α-Synuclein-Y133 Rabbit pAb

ABclonal www.abclonal.com

Catalog No.: AP0451

### **Basic Information**

#### **Observed MW**

20kDa

#### **Calculated MW**

14kDa

### Category

Polyclonal Antibody

# **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse

# **Background**

Alpha-synuclein is a member of the synuclein family, which also includes beta- and gamma-synuclein. Synucleins are abundantly expressed in the brain and alpha- and beta-synuclein inhibit phospholipase D2 selectively. SNCA may serve to integrate presynaptic signaling and membrane trafficking. Defects in SNCA have been implicated in the pathogenesis of Parkinson disease. SNCA peptides are a major component of amyloid plaques in the brains of patients with Alzheimer's disease. Alternatively spliced transcripts encoding different isoforms have been identified for this gene.

# **Recommended Dilutions**

WB

1:500 - 1:2000

### Immunogen Information

Gene ID

**Swiss Prot** 

P37840

#### **Immunogen**

A phospho specific peptide corresponding to residues surrounding Y133 of human alpha-Synuclein

### **Synonyms**

PD1; NACP; PARK1; PARK4; Phospho-α-Synuclein-Y133

### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using Phospho- $\alpha$ -Synuclein-Y133 pAb (AP0451) at 1:1000 dilution.Jurkat and NIH/3T3 cells were treated by Pervanadate (1 mM) at 37°C for 30 minutes. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.