

AP0489

Leader in Biomolecular Solutions for Life Science



# Phospho-YAP1-S127 Rabbit pAb

Catalog No.: AP0489 **29 Publications**

## Basic Information

### Observed MW

75kDa

### Calculated MW

54kDa

### Category

Polyclonal Antibody

### Applications

WB,IHC-P,IF/ICC,ELISA

### Cross-Reactivity

Human,Mouse,Rat

## Background

This gene encodes a downstream nuclear effector of the Hippo signaling pathway which is involved in development, growth, repair, and homeostasis. This gene is known to play a role in the development and progression of multiple cancers as a transcriptional regulator of this signaling pathway and may function as a potential target for cancer treatment. Alternative splicing results in multiple transcript variants encoding different isoforms.

## Recommended Dilutions

|        |               |
|--------|---------------|
| WB     | 1:100 - 1:500 |
| IHC-P  | 1:50 - 1:200  |
| IF/ICC | 1:50 - 1:100  |

## Immunogen Information

### Gene ID

10413

### Swiss Prot

P46937

### Immunogen

A synthetic phosphorylated peptide around S127 of human YAP1 (NP\_001123617.1).

### Synonyms

YAP; YKI; COB1; YAP2; YAP-1; YAP65; Phospho-YAP1-S127

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of lysates from NIH/3T3 cells, using Phospho-YAP1-S127 Rabbit pAb (AP0489) at 1:400 dilution. NIH/3T3 cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 60s.




Immunohistochemistry analysis of Phospho-YAP1-S127 in paraffin-embedded rat lung tissue using Phospho-YAP1-S127 Rabbit pAb (AP0489) at a dilution of 1:25 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of Phospho-YAP1-S127 in paraffin-embedded mouse colon tissue using Phospho-YAP1-S127 Rabbit pAb (AP0489) at a dilution of 1:25 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of Phospho-YAP1-S127 in paraffin-embedded human kidney tissue using Phospho-YAP1-S127 Rabbit pAb (AP0489) at a dilution of 1:25 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of C6 cells (treated with Calyculin A) and C6 cells (untreated) using Phospho-YAP1-S127 Rabbit pAb (AP0489) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.