Leader in Biomolecular Solutions for Life Science

# Phospho-Axl-Y702 Rabbit pAb

Catalog No.: AP0848 1 Publications



### **Basic Information**

### **Observed MW**

138kDa

#### **Calculated MW**

98kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,ELISA

### **Cross-Reactivity**

Human

# **Background**

The protein encoded by this gene is a member of the Tyro3-Axl-Mer (TAM) receptor tyrosine kinase subfamily. The encoded protein possesses an extracellular domain which is composed of two immunoglobulin-like motifs at the N-terminal, followed by two fibronectin type-III motifs. It transduces signals from the extracellular matrix into the cytoplasm by binding to the vitamin K-dependent protein growth arrest-specific 6 (Gas6). This gene may be involved in several cellular functions including growth, migration, aggregation and anti-inflammation in multiple cell types. The encoded protein acts as a host cell receptor for multiple viruses, including Marburg, Ebola and Lassa viruses and is a candidate receptor for the SARS-CoV2 virus.

### **Recommended Dilutions**

**WB** 

1:100 - 1:500

### **Immunogen Information**

Gene ID 558 **Swiss Prot** 

P30530

#### **Immunogen**

A synthetic phosphorylated peptide around Y702 of human AXL (NP\_068713.2).

#### **Synonyms**

ARK; UFO; JTK11; Tyro7; Phospho-Axl-Y702

# Contact

•

www.abclonal.com

### **Product Information**

**Source** Rabbit **Isotype** 

Purification

IgG


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of A-549, using Phospho-Axl-Y702 antibody (AP0848) at 1:400 dilution.A-549 cells were treated by Gas6 (400 ng/ml) at  $37^{\circ}$ C for 5 minutes. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.