

Phospho-PTEN-S380 Rabbit mAb

Catalog No.: AP0998 **Recombinant**

Basic Information

Catalog No.

AP0998

Observed MW

54kDa

Calculated MW

54kDa

Category

Primary antibody

Applications

WB

Cross-Reactivity

Human

Recommended Dilutions

WB 1:500 - 1:2000

Background

This gene was identified as a tumor suppressor that is mutated in a large number of cancers at high frequency. The protein encoded by this gene is a phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase. It contains a tensin like domain as well as a catalytic domain similar to that of the dual specificity protein tyrosine phosphatases. Unlike most of the protein tyrosine phosphatases, this protein preferentially dephosphorylates phosphoinositide substrates. It negatively regulates intracellular levels of phosphatidylinositol-3,4,5-trisphosphate in cells and functions as a tumor suppressor by negatively regulating AKT/PKB signaling pathway. The use of a non-canonical (CUG) upstream initiation site produces a longer isoform that initiates translation with a leucine, and is thought to be preferentially associated with the mitochondrial inner membrane. This longer isoform may help regulate energy metabolism in the mitochondria. A pseudogene of this gene is found on chromosome 9. Alternative splicing and the use of multiple translation start codons results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Feb 2015]

Immunogen Information

Gene ID

5728

Swiss Prot

P60484

Immunogen

A phospho specific peptide corresponding to residues surrounding S380 of human PTEN

Synonyms

10q23del; BZS; CWS1; DEC; GLM2; MHAM; MMAC1; PTEN1; TEP1

Contact

 | www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of 293T cells, using Phospho-PTEN-S380 Rabbit mAb (AP0998) at 1:1000 dilution. 293T cells were treated by CIP(20uL/400ul) at 37°C for 1 hour.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25ug per lane.
Blocking buffer: 3% BSA.
Detection: ECL Basic Kit (RM00020).
Exposure time: 3min.