# Phospho-BRD4-T204 Rabbit pAb

Catalog No.: AP1124



# **Basic Information**

**Observed MW** 200kDa

**Calculated MW** 152kDa

Category **Polyclonal Antibody** 

Applications WB, ELISA

**Cross-Reactivity** Human, Mouse, Rat

# Background

The protein encoded by this gene is homologous to the murine protein MCAP, which associates with chromosomes during mitosis, and to the human RING3 protein, a serine/threonine kinase. Each of these proteins contains two bromodomains, a conserved sequence motif which may be involved in chromatin targeting. This gene has been implicated as the chromosome 19 target of translocation t(15;19)(q13;p13.1), which defines an upper respiratory tract carcinoma in young people. Two alternatively spliced transcript variants have been described.

#### **Recommended Dilutions Immunogen Information**

WB 1:500 - 1:1000 Gene ID Swiss Prot 23476 O60885

#### Immunogen

A synthetic phosphorylated peptide around T204 of human BRD4 (NP\_490597.1).

### **Synonyms**

CAP; MCAP; HUNK1; HUNKI; FSHRG4; Phospho-BRD4-T204

# **Product Information**

Ð www.abclonal.com

Source

Isotype IgG


Purification Affinity purification

## Storage

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

# Validation Data



Western blot analysis of lysates from NIH/3T3 cells, using Phospho-BRD4-T204 Rabbit pAb (AP1124) at 1:1000 dilution.NIH/3T3 cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.