AP1179

Leader in Biomolecular Solutions for Life Science

# Phospho-BRD4-S498/S499 Rabbit pAb

Catalog No.: AP1179



### **Basic Information**

Observed MW 125kDa

Calculated MW 152kDa

Category Polyclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human

### Background

The protein encoded by this gene is homologous to the murine protein MCAP, which associates with chromosomes during mitosis, and to the human RING3 protein, a serine/threonine kinase. Each of these proteins contains two bromodomains, a conserved sequence motif which may be involved in chromatin targeting. This gene has been implicated as the chromosome 19 target of translocation t(15;19)(q13;p13.1), which defines an upper respiratory tract carcinoma in young people. Two alternatively spliced transcript variants have been described.

### **Recommended Dilutions**

### **Immunogen Information**

WB

1:500 - 1:1000

**Gene ID** 23476

Swiss Prot O60885

#### Immunogen

A synthetic phosphorylated peptide around S498 & S499 of human BRD4 (NP\_490597.1).

#### Synonyms

CAP; MCAP; HUNK1; HUNKI; FSHRG4; Phospho-BRD4-S498/S499

Contact

## Product Information


 www.abclonal.com

**Source** Rabbit **Isotype** IgG Purification Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

### **Validation Data**



Western blot analysis of lysates from HeLa cells, using Phospho-BRD4-S498/S499 Rabbit pAb (AP1179) at 1:1000 dilution.HeLa cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serumstarvation overnight. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 180s.