Leader in Biomolecular Solutions for Life Science

# Phospho-GSK3β-S9 Mouse mAb

Catalog No.: AP1258 1 Publications



### **Basic Information**

### **Observed MW**

46kDa

#### **Calculated MW**

47kDa

#### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB,IHC-P,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

### CloneNo number

AMC0523

# **Background**

The protein encoded by this gene is a serine-threonine kinase belonging to the glycogen synthase kinase subfamily. It is a negative regulator of glucose homeostasis and is involved in energy metabolism, inflammation, ER-stress, mitochondrial dysfunction, and apoptotic pathways. Defects in this gene have been associated with Parkinson disease and Alzheimer disease.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

IHC-P 1:200 - 1:500

## **Immunogen Information**

**Gene ID**Swiss Prot
2932
P49841

#### **Immunogen**

A phospho specific peptide corresponding to residues surrounding S9 of human GSK3β

### **Synonyms**

GSK3B; gsk-3β; Phospho-GSK3β-S9

### **Contact**

www.abclonal.com

### **Product Information**


SourceIsotypePurificationMouseIgG1Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.4.

# **Validation Data**



Western blot analysis of various lysates using Phospho-GSK3 $\beta$ -S9 Mouse mAb (AP1258) at 1:1000 dilution.HeLa NIH/3T3 and C6 cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.