

EcoRV



Catalog: RK21106

Size: 4,000 U / 20,000 U

Concentration: 20,000 U/ml

Components:

EcoRV (20,000 U/ml)	RM21604
10X Buffer CutS	RM20103

Product Description

Restriction site:



Unit Definition:

One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Storage Conditions:

10 mM Tris-HCl, 50 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 200 µg/ml BSA, 50% Glycerol, pH 7.4 @ 25°C

Storage Temperature: -20°C

Reaction Conditions:

1X Buffer CutS, Incubate at 37°C.

1X Buffer CutS: 50 mM Potassium Acetate, 20 mM Tris-acetate, 10 mM Magnesium Acetate, 100 µg/ml BSA, pH 7.9 @ 25°C

Quick Cut: Yes

This enzyme digests 1 unit substrate in 5-15 minutes under recommended reaction conditions.

Activity in ABclonal Buffer

CutA	CutB	CutC	CutS
10%	50%	100%	100%

Heat Inactivation: 80°C for 20 min

Methylation Sensitivity:

dam Methylation	Not Sensitive
dcm Methylation	Not Sensitive
CpG Methylation	Impaired by Some Combinations of Overlapping

Instructions

A "Typical" EcoRV Digest:

Composition	Amount
H ₂ O	Up to 50 µl
10X Buffer CutS	5 µl
DNA*	1 µg
EcoRV	0.25-1µl
	10 units are sufficient; generally, 1 µl is used

- ◆ The substrates are completely digested in 5-15 min when incubated at 37°C.
- ◆ *DNA substrates should be free of phenol, chloroform, ethanol, EDTA, detergents, or high concentrations of salt; otherwise, it affects the enzyme activity.

QC Process:

- ◆ Purity is above 95% detected by SDS-PAGE.
- ◆ No exonuclease, nuclease contamination.
- ◆ No residual host genomic DNA detected by PCR.

Optimizing Restriction Endonuclease Reactions

There are several key factors to consider when setting up a restriction endonuclease digest. Using the recommended concentration of DNA, enzyme, and buffer components in the correct reaction volume allows you to achieve optimal digestion. By definition, 1 unit of restriction enzyme completely digests 1 µg of substrate DNA in a 50 µl reaction in 60 minutes. This enzyme : DNA : reaction volume ratio can be used as a guide when designing reactions. However, most researchers follow the "typical" reaction conditions

Order: service@abclonal.com
Tech: support@abclonal.com

For research purposes only. Not for therapeutic or diagnostic purposes in humans or animals.
Please visit <http://abclonal.com> for a complete listing of recommended products.

listed, where a 5–10 fold overdigestion is recommended to overcome variability in DNA source, quantity, and purity.

A "Typical" Restriction Digest

Restriction Enzyme	10 units are sufficient; generally, 1 µl is used
DNA	1 µg
10X ABclonal Buffer	5 µl (1X)
Total Reaction Volume	50 µl
Incubation Time	1 hour*
Incubation Temperature	Enzyme dependent

* Can be decreased to 5-15 minutes by using a "Quick Cut" Restriction Enzyme.

Enzyme

- Keep on ice when not in the freezer.
- Should be the last component added to reaction.
- Mix components by pipetting the reaction mixture up and down or by "flicking" the reaction tube. Follow with a quick ("touch") spin-down in a microcentrifuge. Do not vortex the reaction.
- In general, we recommend 5–10 units of enzyme per µg DNA, and 10–20 units for genomic DNA in a 1 hour digest.

DNA

- Should be free of contaminants such as phenol, chloroform, alcohol, EDTA, detergents, or excessive salts. Extra wash steps during purification are recommended.
- Methylation of DNA can inhibit digestion with certain enzymes.

Buffer

- Use at a 1X concentration.
- Supplement with SAM (S-Adenosyl methionine) to the recommended concentration if required.

Reaction Volume

- A 50 µl reaction volume is recommended for digestion of 1 µg of substrate.
- Enzyme volume should not exceed 10% of the total reaction volume to prevent star activity due to excess glycerol.
- Additives in the restriction enzyme storage buffer (e.g.,

glycerol, salt), as well as contaminants found in the substrate solution (e.g., salt, EDTA, or alcohol), can be problematic in smaller reaction volumes. The following guidelines can be used for techniques that require smaller reaction volumes.

Reaction System	Enzyme Amount*	DNA	10X ABclonal Buffer
10 µl**	1 U	0.1 µg	1 µl
25 µl	5 U	0.5 µg	2.5 µl
50 µl	10 U	1 µg	5 µl

* Restriction Enzymes should be diluted when smaller amounts are needed. ** 10 µl rxns should not be incubated for longer than 1 hour to avoid evaporation.

Incubation Time

- Incubation time is typically 1 hour.
- Can often be decreased by using an excess of enzyme, or by using one of our "Quick Cut" restriction enzymes.
- It is possible, with many enzymes, to use fewer units and digest for up to 16 hours.

Stopping a Reaction

If no further manipulation of DNA is required:

- Terminate with a stop solution (10 µl per 50 µl rxn) [1x: 2.5% FicoII-400, 10 mM EDTA, 3.3 mM Tris-HCl, 0.08% SDS, 0.02% Tartrazine, 0.001% Xylene Cyanol FF, pH 8.0 @ 25°C].

When further manipulation of DNA is required:

- Heat inactivation can be used.
- Remove enzyme by using a spin column or phenol/chloroform extraction

Control Reactions

If you are having difficulty cleaving your DNA substrate, we recommend the following control reactions:

- Control DNA (DNA with multiple known sites for the enzyme, e.g., lambda or pUC19 DNA) with restriction enzyme to test enzyme viability.

If the control DNA is cleaved and the experimental DNA resists cleavage, the two DNAs can be mixed to determine if an inhibitor is present in the experimental sample. If an inhibitor (often salt, EDTA, or phenol) is present, the control DNA will not cut after mixing.

Order: service@abclonal.com
Tech: support@abclonal.com

For research purposes only. Not for therapeutic or diagnostic purposes in humans or animals.
Please visit <http://abclonal.com> for a complete listing of recommended products.