

T4 DNA Ligase



Catalog: RK21501

Size: 16,000 U / 80,000 U

Concentration: 400,000 U/ml

Components:

T4 DNA Ligase (400,000 U/ml) RM21501

10X T4 DNA ligase Reaction Buffer RM20108

Product Description

T4 DNA Ligase can catalyze the formation of a phosphodiester bond between juxtaposed 5' phosphate and 3' hydroxyl termini in duplex DNA or RNA. This enzyme joins blunt-end and cohesive-end termini as well as repairs single-stranded nicks in duplex DNA and some DNA/RNA hybrids. T4 DNA Ligase seals nicks for these DNA substrates. T4 DNA Ligase is applicable to cloning restriction fragments and to joining linkers and adapters to blunt-ended DNA.

Product Source: An *E. coli* strain that carries the T4 DNA ligase gene.

Unit Definition:

One unit is defined as the amount of enzyme required to give 50% ligation of HindIII fragments and 50% of HindIII digestion fragments of λ DNA (5' DNA termini concentration of 0.12 μ M, 300 μ g/ml) in a total reaction volume of 20 μ l over 30 minutes at 16°C in 1X T4 DNA Ligase Reaction Buffer.

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, pH7.4 @ 25°C

Storage Temperature: -20°C

Reaction Conditions:

1X T4 DNA Ligase Reaction Buffer.

1X T4 DNA Ligase Reaction Buffer: 50 mM Tris-HCl, 10 mM MgCl₂, 10 mM DTT, 1 mM ATP, pH7.5 @ 25°C

Heat Inactivation: 65°C for 10 min.

Instructions

- ◆ Set up the following reaction in a microcentrifuge tube on ice.

Composition	Amount
10X T4 DNA Ligase Reaction Buffer*	2 μ l
Vector DNA (4 kb)	50 ng (0.02 pmol)
Insert DNA (1 kb)**	37.5 ng (0.06 pmol)
Nuclease-free dH ₂ O	up to 19 μ l
T4 DNA Ligase ***	1 μ l
Volume	20 μ l

*:10X T4 DNA Ligase Reaction Buffer should be thawed and resuspended at room temperature.

** Insert DNA (1 kb): A ligation using a molar ratio of 1:3 vector to insert for the indicated DNA sizes.

***:T4 DNA Ligase should be added last.

- ◆ Short centrifugation after gentle percussion.
- ◆ Gently mix the reaction by pipetting up and down and microfuge briefly.
- ◆ For cohesive (sticky) ends, incubate at 16°C overnight or room temperature for 10 minutes.
- ◆ For blunt-ends or single-base overhangs, incubate at 16°C overnight or room temperature for 2 hours (alternatively, high-concentration T4 DNA Ligase can be used in a 10-minute ligation).
- ◆ Heat inactivate at 65°C for 10 minutes.
- ◆ Chill on ice and transform 1-5 μ l of the reaction into 50 μ l competent cells.

QC Process:

- Purity is above 95% detected by SDS-PAGE.
- No exonuclease, nuclease, RNase contamination.
- No residual host genomic DNA detected by PCR.

Order: service@abclonal.com
Tech: support@abclonal.com

For research purposes only. Not for therapeutic or diagnostic purposes in humans or animals.
Please visit <http://abclonal.com> for a complete listing of recommended products.